Missouri Botanical Garden

Peter H. Raven Library

George Engelmann Botanical Notebooks



Pagination Note:
Since many of the items lack a specific page number, the page number displayed online refers to the sequentially created number each item was given upon cataloging the materials.

39 05 h.s/o

L. ~~petiolata~~ serrula (petals 1.)

annua decumbens vale
patenter pilosa

~~ploribus~~ foliis ebasi valde obliqua
obtuse-ovata v. subrotunda.
oblonga ~~ovata~~ basi obtusis obtusis
acute ~~serratis~~ grosse pauci serratis
subtus pilosis supra subnudis
stipulis lanceolatis carinatis
glanduli catenulibus
appendiculibus auguste obtegens
ovatulatis

~~Ovaria~~ Styli ~~bacillus~~
~~petiolatis~~ ovario globoso brevissimo
pilulis adnato ferè basi libero
capsula acute angulata
Semina ovata ~~acute~~ laevia.
~~acute compressa~~ obtuse angulata

Bracteæ 4—6 subulatæ
laeves very sharply serrate
2—5 l. lg 1—3 broad
Leaves ~~Dec~~ lg —0.6 l. & dentate
remarkably ~~auric~~ winged angles

Found in New Mexico St Fe.

№ 796 ÷ 804
Wright 1843
1844 (547 1852) Escondido Creek west of El Paso
(304. 1851) Pecos
(817 (1849) № 658 west of Pecos ÷ Torrey
(59. 1851 ÷ part) S Diana
(796 1851)

6628

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

29/34

Euph. villifera Scheele

annua erecta patula ramosissima
villis patulis hirsuta

foliis ellipticis vel obliqueovatis subcordatis ovatis obliquis integris vel
grosse dentatis, utrinque pilosis
stipulis subulatis deciduis
auctorum alarum
apice ramulorum corymbosis
involucris globosis, appendiculis
obovatis patulis (albis v. roseis)
stylis patulis ovario globoso
brevioribus ultra medio 6-fidis.
capsula oblongo angulata
seminibus globoso ovatis obtuse
angulatis, subtuberculatis.

Euphorbia villifera, Cedarwood on rocky hill,
No Burkes R.L. No 530
Grey. 350. near Salsillo
Berlandier 2084 & 674
Wright.

A bushy branched plant 6-12 inches
high; leaves and shoots thickly
clothed with spreading white
hairs; stems and branches usually
by a thick, dense, & fleshy
spreading thickened base;
3-4 lines long 2-3 broad; of a
dark green red — seeds about 0.5 l. long.
and nearly 0.4 thick. — Bracts
opposite are more naked
and have scarcely dentate
leaves. So are the next two
ones from Salsillo, they grow more
are only half a foot or a few inches
high, have very much compressed
capsules & smaller seeds.

Seems to be near E. velligera Schauer
in Linn XX. 728 — but that has villose capsules —

or

E. velligera
Schauer
Wish maybe
the same, but
has serrulate leaves
& villose
capsules

6629

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Among Coulters' Plants 1447 is an Euphorbia (mixed with 2 others) which I have not classed:

Annua? caule decumbente ramoso pilosis e basi obliqua obtuse subcordata ovatis seu obovatis integerrimis dein crenulatis subtus pilosis; stipulis lanceolatis floribus parcis involucris glandulis 4 auguste appendiculatis. stylis adnexis bifidis; capsulis obtuse angulatis medio sulcatis (0.5 li) ante angulatos pruinoso rugosis

perhaps mere E. pilosula??



cm 0 1 2 3 4 5 6 7 8 9 10
copyright reserved
MISSOURI BOTANICAL GARDEN

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

6630

Or (b) 5 (p)

(2 pag 5

Euphorbia hyssopifolia

annua caule erecto seu diffuso
puberulo seu glabrato

foliis e basi obliqua oblonga
seu subovata ovato-oblongis
s.s. oblongo-lanceolatis falcatisve
obtusis argute serratis subtus
glabris pallidis supra pilosis
pilosis seu nudis saepe punc-
tatis

stipulis ebasi triangulato
lanceolatis fimbriatis.

anthodiis alaribus demum
ad ramulos alternatim
apice in cymulas laxe
foliosas collectis;

appendiculibus parvis
orbiculatis integris, seu raro
subnullis.

stylis glabris profunde bifidis, ramis
subulatis divaricatis.

capsula glabra, obtuse angula
seminibus ovatis obtuse
angulatis rugulosis tuberculatis
nigricantibus.

var. brevifolia see below

Var. cymulis laeta (see No 4 pag 5)

suffruticosa glaberrima
foliis confertis lanceolatis acutis serrulatis acutis
cappadicibus lactis orbiculatis
candidis stylis basi coalitis
cymulis laxifloris longe pedunculatis nudis
demum in paniculam terminalem dispositis

6631

Berlandier 489
& 1849

Charpres Fendler 277

Stigmatibus clavellatis.

Common throughout the U. S. on moist soil, in fields & waste places

Var. Crocifolia Diffusa pilosa foliis ovatis s. inferioribus subcordatis pectinato-dentatis calycibus vix ac capsulas longioribus.

Texas Lindheimer. (Devil River)
New Mexico Wright 1842 in part
Bigelow 655 (1849. 882)
Fendler
Mexico Gregg 137 Hartweg 32 Coult 1436
Guyana Schomb Gray
Chili
Panama Fendler 277
Cuba Thurber 1065
Upper Missouri Hayden
Var Brevifolia — N. castern & Northwestern states, & the Rocky Mts. — but Mexico & California
Magalia Coll by S. Gregg

Annual Known weed
usually 1–2 feet high
or long — rarely 2, rar 3
often 4 inches high —
leaves very variable
broadly ovate — to long
linear, usually 10–18
lines long, 3–6 or even 8 lines
broad. 0,25 — 0,6 lines
long, very thick & the
only stalk (long) in the
single species! —
Var Cuneata from Mexico.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

~~40~~ 25

It seems difficult to consider that this should be the original limen plant, but as it grows it is even when known and acknowledged under this name it seems impossible to make any changes, before having examined them from Cabruca. The original plant seems to be some undescribed species.

It is sufficiently distinct, entirely glabrous, with large petal like appendages, different leaves, but undistinguishable seeds. It is perhaps H. & B. C. pulchella, but this is said to have almost entire leaves.

E. dumiflora is probably only a form! The Key west plant is distinct enough but outlines forms unite it with Euph. hyssopifolia.

Tuerta, Venezuela 1186 is a peculiar variety with very pubescent capsules! Whole plant more hairy than usual; one specimen has bract appendages, the other almost none!

6632

Euphorbia Hubrica n. sp.

E. hypericifolia Planta Kotschyana.
non Lin

annua (?) robusta erecta s.

decumbens;

foliis ex basi valde obliquis
ovatis s. oblongis, obtus. mucronulatis
vix ciliatis s. parce s. obtusiuscule
serrulatis

stipulis triangulares acutis,
undulatis

floralibus ovatis, ovato-lanceolatis
s. lanceolatis, subfalcatis, subcordatis, truncatis
appendiculatis
capsulis pubescentibus.

Kotschy 334

α. glabra Kotschy Pl. Nub. 154

β. pubescens

glabra, foliis oblongis lanceolatis
capsulis glabris

β. pubescens tota pubescens
foliis ovatis lanceolatis
capsulis pubescentibus. Kotschy. 334

Differs from Eupt. hyperic. in the pale narrow shapeless seeds — the almost entire leaves — the absence of any obtuseish — style short.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

Eupt. Prolii Guss. fide spec. in hb Guss. ex Palermo
is somewhat inappendiculate form of the common E. hyperic.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

#95.6 E. pauciflora — OT 95p

probably a form of E. hyperi-

E. densiflora

annua glabra erecta glabra

foliis e basi oblongo oblanceolatis

argute serratis,

stipulis saepe connatis

truncatis, ut ergo ramulis

~~foliosis~~ ramis floriferis

alternis apice glomerulum multiflorum

basi foliis oppositis ~~foliosum~~ ~~foliosum~~ & subaphyllum

gerentibus ~~foliosum~~

~~basi~~ appendiculis ~~ovatis~~

angustis integris & nullis

stylis brevibus ad basi

capsula globosa

seminibus ovatis brunneis

reticulosis subcellis s. foveis

Key West, Florida, Chapman, Blodgett

Monterey, Mexico Dr Edwards

Chagres Pl. Coll. 275

Venezuela " " 1185 & 1187

Porto Rico

Very similar in appearance

to E. hyperifolia, and perhaps

the ~~true~~ typical Linn plant

~~of that name~~.

Leaves 3/4—1 1/2 inches long,

1/4—6 lines wide — always smooth

Glomerules little large ± 10 fl.

In S. Amer & Mexico

more copiously — than in

the West Indies & Key West

3—6 lines ± Santa ± in a Venezuelan

Specimen 2 fl. to each axil

6633

and in any Botrus more or small or unusually large ones in the same head, dissimilar objects I have made with on E. cephalantha. Leeds similar in shape to those of E. hypericifolia but only 0.4 C. long

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

42
96

ax

psp

x pg 1.

E. pilulifera L.

annua, erecta s. patula, valde
pilis flavis articulatis tecta
s. adnodos solum hirsuta,
foliis e basi oblique
acuta s. mucronulata obtusiusve
basi cuneatis; obtuse
serratis utrinque [illegible]
stipulis persistentibus
stipulis subulatis
glomeris globosis dense pauci-
multifloris, pseudoaxillaribus alternis
sessilibus seu pedunculatis.
glomeris involucrum multi-
stipitatum
appendiculatis
stylis apice s. ad medium
bipartitis dense hispidulo
papillosis; stigmate globoso capitellato
semiglobo
acute angulatus transverse
rugoso rosei

β. *E. viatorum*.
γ. *xylorhiza*

Keyserl. Blodgett
Mexico Coulter 1438 ÷ part & γ. Gray
Chihuahua Pringle 274 ÷ Wright
China Hale Gray. Mexican Boundary
Wright 1842 ÷ part (535 1852 2d) Sonora

Upland prairies, rocky
ravines, &c. in the
Globose involucrate heads

6634

Spreading 6— 15 inches high —
leaves 1½— 1½ inches long
always acute at base. — Florida
Apalachicola Bay — Yellow
with obtuse. at tip — Yellow
Pericarp thin always present
& quite persistent — always having
T. Capsule 0.4 by 0.3 seeds
only 0.3 by — lengthwise
Marked in the Glumes,
Key West Spruce on Bark
Jubula appendages, (all the others
very large & conspicuous appendages.
Among Collins 1438 is much
allied & perhaps well-distinguished
for which seems to constitute
one species. —

Localities for E. pilulifera

Rio Janeiro (Hb Bram.
Martinique "
Isle de France "

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN HERBARIUM

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

4426/5.

E. d'Urville

annua adscendens seu diffusa
pilosa
foliis e basi obliqua
ovatis v. lanceolatis subrhombeis
acutiusculis serratis
stipulis subulatis
glomerulis globosis pseudo-
axillaribus sessilibus seu
breviter pedunculatis aphyllis
involucris minutis pubendis
glandulis stipitatis, appendi-
culatis paucis
stylis ad medium bifidis
brevibus; stigmatibus
capitatis
; capsula pilosa
semine minutè ovato
acutè rugoso tuberculato.

Wright 1892 in part
(535. 1851 2d) Sonora rock pebbly bed

Stems ascending ½–1 ft long
1–1½ inches in length. ¼–½ inch
in breadth. with a lanceolate
red spot above the middle the
very pale or glaucous the lower
side (Involucres hemispherical
½ line long, very much crowded
heads 3–4 lines in diameter
capsules very small shining
seeds below 0,3 to 0.4 line in
length.

Nearly allied to E. pilulifera

6635

MISSOURI BOTANICAL GARDEN
ST. LOUIS ENGELMANN HERBARIA

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

~~15 37~~

~~cymulosa~~

Euph. ~~cephalanthus~~ capitellata

esotrichoides s. dem.

annua diffusa caulibus elongatis

pubentibus

foliis e basi valde obliqua

oblongo linearibus nudis seu parce

~~hirtellis~~ (micronatis) argute

serrulatis

stipulis lanceolatis subulato —

laceris

glomeris demifloris semialtis fere

~~capitellatis~~ aphyllis

Involucri glandulis ~~obtectulato~~

~~appendiculatis~~ elongatis stipitatis ~~obovato~~ appendi-

culatis v. late obovatis

integris

styliis ovario hirto longioribus

~~filiformibus~~ ad med — bifidis

elongatis

capsula pubescente

dentibus ovatis acutatis

~~fusco~~ angulatis rugoso

tuberculosis.

Wright 1849 (~~1851 & 99~~) Cow valley

San Bernardino, Sonora

~~Arizona~~ Stems 6—10

inches long; leaves 5—8 lines long

2 l. wide — very sharply serrate;

heads of (anthodia flowering like)

rather showy 4—6 lines

diameter; seed = 0.5 l. long

— Very similar to E. pycnantha

but easily distinguished by the straggling

habit (the naked glomerules

the narrower sharply serrate leaves and the

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

6636

seeds at Strasburg Stable

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

~~44~~ 38

Euphorbia pycnanthema

See Euph. Buttoniana

perennis erecto patula ramosa-
pubescens
foliis ellipticis obliquis
cordato ovatis obtusis s. obtusiusculis
saepe mucronulatis integris
sub dentibus paucis ad ~~basin~~ margines
inferiorem serrulatis subtus
stipulis lanceolato-subulatis
fissis
glomerulis multifloris
terminalibus foliis superantibus plurib
quasi involucratis
involucris pilosulis ~~extrorsis~~
late obovatis appendiculatis
sem (intensibus? subundis)
stylis fere adbasi fissis
patulis stigmatibus divaricatis
clavellatis
capsula subglobosa puberula
semin. ovatis acute angulatis
transverse rugulosis punctatisque

Wright (186.) Mountain sides near Lake Santa Maria

Many stems 6 - 12 inches high from
a very thick woody root
Leaves 4-5 lines long - Seeds
0.5 line long. - ~~Involucre~~ Very much
like a specimen of a small tuberous plant
with a terminal involucrate head!

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

MISSOURI BOTANICAL GARDEN
copyright reserved
cm 0 1 2 3 4 5 6 7 8 9 10

25 ? 105/p.

Euphorbia pubescula

annua? decumbens caescens,
pubescula
folii e basi maxime obl. ovati
subcordata ovati obtusi
crenati introrsum acquitibus seu longioribus
stipulis subulatis fimbriatis
Dutta involucris hemisphaericis
pubescentibus, appendicibus ovatis
patentibus multo majoribus
stylis bipartitis apice
bifidis, ovario caescente
papillatibus longioribus.

Key west, Florida, Mr Blodgett.
small species clearly abt
allied to ? almost imbricate
— Approaches E. dentata in habit
the peculiarly unequal
size of the glands and especially
the petaloid appendages.

GEORGE ENGELMANN PAPERS
MISSOURI BOTANICAL GARDEN

6637



0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

6638

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS



~~25~~ 40 Cata

Euphorbia d'Catata Torr. ~~& Hook~~ & Gray = Pepis Torr. Repens non Hochstetter — Rich. Flor. Abyss

perennis patula seu diffusa humilis rigida pilis adpressis
obsessa

foliis e basi d'Catata obtusa angustata falcatis obtusis margine integerrimo revolutis supra sabris subtus (cum costa utrinque stipulis subulatis deciduis

anthodiis versus apicem caulis singulis s. ~~stipulis~~ = glomerulos pauciflores laxiusculos confertis

involucris ovatis appendiculatis angustis crenatis seu nullis

stylis patulis recurvisque ovar. — caruncula aequantibus seu superantibus ad medium bilobis stigmatibus clavellatis

capsula cum ovulo

angulata

seminibus angulato ovatis
acute angulatis laeviter
tuberculatis.

Wright. 1841 (1846. 1854) San Pedro, Lowry Pass Expedition

Gravelly hills, Eagle Springs, Arizona

Sandy prairies on the Cimaron, Feb 1st 794

A low plant a span high spreading; leaves 1/3 — 1/2 inch long, ~~& roughish~~ half as wide, of a peculiar & characteristic shape, which distinguishes this species at once from the allied ones in the Pedler Specimen the leaves

6639

the lowers
are shorter & broader —
proportion. —

Leaf 0.8 l. by — half as
thick.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

~~24~~ 41 OS pej

E. acuta

~~pr~~ perennis ~~multiculis~~ culmis foliosa hirsuta s.
glaucescens
foliis obtusa ~~obtusis~~ obtusis ubique
lanceolatis, ~~asperis~~ acutis, ~~rigide~~ costulatis
nervosis. supra ~~glabris~~ nudis
stipulis filiformibus mox
deciduis
antheris ovato oblongis apice
clausis (~~3-angulis~~ s. pseudoquadrangulis) appendiculatis
involucris hemisphaericis
appendiculatis hirsutis ~~decaphyllis~~, foliolis
ovatis ~~oblongis~~ obtusis, ~~alternis~~ medio bilobis
petalis / ovario ca no longioribus
capsula cana
seminibus subglobosis (rubris) acute
angulatis laevibus.

Wright (annal 355)
(— — 843)
626)
Sabinal, west bank Rio Frio
west of Reos
San Felipe near
N. 1840 (92, 1851) east of Reos
On the same Reos
1739 (oct to dec 1839)
(~~prob 39 1851~~) Rio Frio
from the Rio Grande to the Lago
(~~355 prob~~) Sabinal
N. 557. (1849) ?
Boyleau on the Reos.

½ – 1 foot high, stiff rigid, rough hairy,
leaves 6–10 lines long 3–6 wide
well channeled by the very
acute point. – Seeds 1 line or
over in length, 0.6 – 0.7 l. thick

6640

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN